
# CASSINI



# TITAN 037TI(T23) MISSION DESCRIPTION


January 2007

**Jet Propulsion Laboratory**California Institute of Technology

#### 1.0 OVERVIEW

After just 16-days from its last flyby of Titan, Cassini returns for its twenty-fourth targeted encounter, Titan-23. The closest approach to Titan occurs on Saturday, January 13, at spacecraft time 2007-013T08:38:31 (Dec. 12 at 3:41AM Pacific Time) at an altitude of 1000 kilometers (~621 miles) above the surface and at a speed of 6.0 kilometers per second (13,400 mph). The latitude at closest approach is 31degrees N (over an area known as Aaru), and the encounter occurs on orbit number 37.

This encounter is set up with two maneuvers: an apoapsis maneuver on January 5, and an approach maneuver, scheduled for January 10. This inbound encounter occurs about 3 days before Saturn closest approach.



#### 1.1 ABOUT TITAN

If Titan were a planet, it would likely stand out as the most important planet in the solar system for humans to explore. Titan, the size of a terrestrial planet, has a dense atmosphere of nitrogen and methane and a surface covered with organic material. It is Titan that is arguably Earth's sister world and the Cassini-Huygens mission considers Titan among its highest priorities.

Although it is far colder and lacks liquid water, the chemical composition of Titan's atmosphere resembles that of early Earth. This, along with the organic chemistry that takes place in Titan's atmosphere, prompts scientists to believe that Titan could provide a laboratory for seeking insight into the origins of life on Earth. Data from the Huygens probe, which touched down on Titan's surface in January 2005, and the Cassini orbiter has shown that many of the processes that occur on Earth also apparently take place on Titan – wind, rain, volcanism, tectonic activity, as well as river channels, and drainage patterns all seem to contribute in shaping Titan's surface. However, at an inhospitable - 290 degrees Fahrenheit (-179 degrees Celsius), the chemistry that drives these processes is fundamentally different from Earth's. For example, it is methane that performs many of the same functions on Titan that water does on Earth.

The Huygens probe landed near a bright region now called Adiri, and photographed light hills with dark river beds that empty into a dark plain. It was believed that this dark plain could be a lake or at least a muddy material, but it is now known that Huygens landed in the dark region, and it is solid. Scientists believe it only rains occasionally on Titan, but the rains are extremely fierce when they come.

Only a small number of impact craters have been discovered. This suggests that Titan's surface is constantly being resurfaced by a fluid mixture of water and possibly ammonia, believed to be expelled from volcanoes and hot springs. Some surface features, such as lobate flows, appear to be volcanic structures. Volcanism is now believed to be a significant source of methane in Titan's atmosphere. However, there are no oceans of hydrocarbons as previously hypothesized. Dunes cover large areas of the surface.

The Cassini-Huygens mission, using wavelengths ranging from ultraviolet to radio, is methodically and consistently revealing Titan and answering long-held questions regarding Titan's interior, surface, atmosphere, and the complex interaction with Saturn's magnetosphere. While many pieces of the puzzle are yet to be found, with each Titan flyby comes a new data set that furthers our understanding of this world as we attempt to constrain scenarios for the formation and evolution of Titan and its atmosphere.


#### 1.2 TITAN-23 SCIENCE HIGHLIGHTS

– RADAR: T23 Synthetic Aperture RADAR (SAR) measurement clips Ganesa Macula, which is thought to be a cryovolcanic dome formed when ammonia and water seeped out onto the surface of Titan. (There may be some topographic constraints, certainly in the imaging of northern, as-yet-unseen part of Ganesa).

This flyby also overlaps with TA SAR and TA/T3 altimetry. End of swath is close to mountains

seen by VIMS and 'rim' of circular feature; T23 also has the highest-latitude altimetry in the mission so far.

- UVIS has an FUV observation, observing the star Eta Ursa Major as it becomes occulted by Titan. See picture to right.
- The ISS outbound and inbound observations will image the surface of Titan and attempt to track atmospheric and surface changes. ISS will also be looking at surface color variations and monitoring limb hazes as well as particle properties, including vertical distributions with resolutions of ~6 km/pix.




- MAPS Analysis of plasma wake, ions escaping from Titan, and Titan's interaction with Saturn's magnetosphere down to a low altitude (1000 kilometers). INMS specifically will be looking at the atmospheric and ionospheric thermal structure of Titan's atmosphere.
- MIMI will investigate micro-scale and near aspects of the Titan interaction by observing during about one hour period around the Titan encounter. With -Y pointed toward Titan, when within 30 minutes of the targeted flyby, it will also measure Titan exosphere/ magnetosphere interaction.

#### 1.3 SAMPLE SNAPSHOTS


Three views of Titan from Cassini before, during, and after closest approach to Titan are shown below. The views are oriented such that the direction towards the top of the page is aligned with the Titan North Pole. The optical fields of view are shown in the first two plots and the remote sensing instrument fields of view are shown in the third, assuming they are pointed towards the center of Titan. The sizes of these fields of view vary as a function of the distance between Cassini and Titan. A key for use in identifying the remote sensing instruments fields of view in the figures is listed at the top of the next page.

Key to ORS Instrument Fields of View in Figures


Instrument Field of View	Depiction in Figure		
ISS WAC (imaging wide angle camera)	Largest square		
VIMS (visual and infrared mapping spectrometer)	Next largest pink square		
ISS NAC (imaging narrow angle camera)	Smallest green square		
CIRS (composite infrared spectrometer) – Focal Plane 1	Small red circle near ISS_NAC FOV		
UVIS (ultraviolet imaging spectrometer)	Vertical purple rectangle centered		
	within largest square		



View of Titan from Cassini two hours before Titan-23 closest approach



View of Titan from Cassini at Titan-23 Closest Approach



View of Titan from Cassini two hours after Titan-23 closest approach

# Timeline and Geometry Table below

## Cassini Titan-23 Timeline - January 2007

Colors: yellow = maneuvers; blue = geometry; pink = T23-related; green = data playbacks

targeting maneuver  ted to accommodate flyby time in on the thermal in on the thermal in on the thermal in on the thermal in stratosphere in on the thermal in stratosphere itside Exobase to lso look for trace an's stratosphere.
targeting maneuver  sed to accommodate flyby time n on the thermal n's stratosphere ide for Atmospheric and Auras n on the thermal n's stratosphere attside Exobase to lso look for trace an's stratosphere.  DAR Radiometry
med to accommodate flyby time n on the thermal n's stratosphere ide for Atmospheric and Auras n on the thermal n's stratosphere attside Exobase to lso look for trace an's stratosphere.  ADAR Radiometry
n on the thermal n's stratosphere ide for Atmospheric and Auras n on the thermal n's stratosphere atside Exobase to lso look for trace an's stratosphere.
n on the thermal n's stratosphere ide for Atmospheric and Auras n on the thermal n's stratosphere atside Exobase to lso look for trace an's stratosphere.
n on the thermal n's stratosphere ide for Atmospheric and Auras n on the thermal n's stratosphere atside Exobase to lso look for trace an's stratosphere.
n on the thermal n's stratosphere ide for Atmospheric and Auras n on the thermal n's stratosphere atside Exobase to lso look for trace an's stratosphere.
n's stratosphere ide for Atmospheric ide Auras n on the thermal n's stratosphere utside Exobase to lso look for trace an's stratosphere.  DAR Radiometry
nnd Auras n on the thermal n's stratosphere atside Exobase to also look for trace an's stratosphere.  DAR Radiometry
n's stratosphere  utside Exobase to lso look for trace an's stratosphere.  NDAR Radiometry
lso look for trace an's stratosphere.
AR Scatterometry tions
to compensate for corque and target pensation
of Titan's surface ing Nadir.
ace features. T23 r lakes on Titan at ern Latitudes.
I miles) , C/A Time, , Speed = 6.0 km/s = Lit outbound and Local Saturn Time = , Magnetospheric Subspacecraft Lat, 1, 358
Synthetic Aperture of Titan's Surface
e of Titan Surface s pointing Nadir
DAR Scatterometry
greater spacecraft ity.
ADAR Radiometry
ons for surface and c mapping
sed to accommodate flyby time
to Goldstone 70M
-51 deg, phase = 39

OWLT (mins)	69.26						
C/A Time		Sat	Jan	13	01:48	AM	

## 1.4 FLYBY GEOMETRY

The Flyby Geometry table is not available for Titan-23.

# 1.5 PLAYBACK TIMELINE

The Playback Timeline is not available for Titan-23